



Otezla® (apremilast) is a prescription medicine used to treat adult patients with:

- Plaque psoriasis for whom phototherapy or systemic therapy is appropriate.
- Active psoriatic arthritis.

Recently prescribed Otezla?

USE THIS TO GET STARTED

Click the links below for information you may need as you get started with treatment.



Commercially insured patients can get Otezla for \$0 per month with the Otezla Co-Pay* Card

If you are commercially insured, you can get Otezla for **\$0 each month**.

How to enroll:

- Go to otezla.com/copay
- Call **1-844-4OTEZLA** (1-844-468-3952)



Helpful resources when getting started

See videos at links below for what to expect when starting treatment.

If you're taking Otezla for plaque psoriasis:

What results might I expect and when? [Watch video](#)

What are the common side effects? [Watch video](#)

If you're taking Otezla for psoriatic arthritis:

What results might I expect and when? [Watch video](#)

What are the common side effects? [Watch video](#)

You can also check out otezla.com or download a digital brochure specifically for getting started with treatment at the button below.

DOWNLOAD BROCHURE 



What's next?

Your Otezla prescription will be delivered right to your doorstep by your specialty pharmacy.

[Click here](#) to learn more.

Have questions along the way? We're here for you.

Our team is available 8 AM – 8 PM ET, Monday – Friday at **1-844-4OTEZLA** (1-844-468-3952).

*The Otezla Co-Pay Card is for eligible commercially insured patients. Terms, conditions, and program maximums apply. Other restrictions may apply. Please see full Terms and Conditions at otezla.com/copay. This program is not open to patients receiving prescription reimbursement under any federal, state, or government-funded healthcare program. Not valid where prohibited by law. Patients who are not eligible can call 1-844-4OTEZLA to discuss other financial support opportunities.

IMPORTANT SAFETY INFORMATION

You must not take Otezla if you are allergic to apremilast or to any of the ingredients in Otezla.

Otezla can cause allergic reactions, sometimes severe. Stop using Otezla and call your healthcare provider or seek emergency help right away if you develop any of the following symptoms of a serious allergic reaction: trouble breathing or swallowing, raised bumps (hives), rash or itching, swelling of the face, lips, tongue, throat or arms.

Please see additional Important Safety Information continued on back and Full Prescribing Information [here](#).

IMPORTANT SAFETY INFORMATION (continued)

Otezla can cause severe diarrhea, nausea, and vomiting, especially within the first few weeks of treatment. Use in elderly patients and the use of certain medications with Otezla appears to increase the risk of complications from having severe diarrhea, nausea, or vomiting. Tell your doctor if any of these conditions occur.

Otezla is associated with an increase in depression. In clinical studies, some patients reported depression, or suicidal behavior while taking Otezla. Some patients stopped taking Otezla due to depression. Before starting Otezla, tell your doctor if you have had feelings of depression, or suicidal thoughts or behavior. Be sure to tell your doctor if any of these symptoms or other mood changes develop or worsen during treatment with Otezla.

Some patients taking Otezla lost body weight. Your doctor should monitor your weight regularly. If unexplained or significant weight loss occurs, your doctor will decide if you should continue taking Otezla.

Some medicines may make Otezla less effective and should not be taken with Otezla. Tell your doctor about all the medicines you take, including prescription and nonprescription medicines.

The most common side effects of Otezla include diarrhea, nausea, upper respiratory tract infection, tension headache, and headache. These are not all the possible side effects with Otezla. Ask your doctor about other potential side effects. Tell your doctor about any side effect that bothers you or does not go away.

Tell your doctor if you are pregnant, planning to become pregnant or planning to breastfeed.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call **1-800-332-1088**.

Please [click here](#) for the Full Prescribing Information.